


Agromet Advisory Bulletin for Feni District

		
	 Agro-Meteorological Information Systems Development Project Component-C of BWCSR Department of Agricultural Extension	
Date: 24th June 2020 Bulletin No. 157	Agromet Advisory Bulletin for Feni District (24th June to 28th June 2020)	

Weather Conditions for last four days (20th June to 23rd June 2020)

Weather Parameters	20.06.20	21.06.20	22.06.20	23.06.20	Range
Rainfall (mm)	1.0	5.0	5.0	23.0	1.0-23.0 (34.0)
Maximum Temperature (° C)	30.7	32.8	32.8	33.0	30.7-33.0
Minimum Temperature (° C)	25.4	26.0	25.5	25.5	25.4-26.0
Relative Humidity (%)	65.0-93.0	71.0-96.0	63.0-93.0	77.0-95.0	63-96
Wind Speed (km/h)	5.6	7.4	9.2	3.7	3.7-9.25
Cloud Amount (Okta)	5	8	7	7	5-8
Wind Direction	South/South-westerly	South/South-westerly	South/South-westerly	South/South-westerly	South/South-westerly

Weather forecast as per Bangladesh Meteorological Department for the next 5 days (24th June to 28th June 2020)

Weather Parameters	Range
Rainfall (mm)	0.0-2.2 (5.4)
Maximum Temperature (° C)	30.1-31.1
Minimum Temperature (° C)	26.2-26.9
Relative Humidity (%)	82.0-88.0
Wind Speed (Km/h)	4.8-7.6
Cloud status	Partly Cloudy Sky
Wind Direction	South/South-westerly

Agromet Advisories

Special Agromet Advisories for CORONA virus (COVID-19) transmission in Bangladesh

Farmers and all other farm managements are advised to maintain the social distancing (at least 3 meters), cover their faces with masks and follow other guidelines of Government of Bangladesh to avoid infection and social transmission of CORONA virus (COVID-19).

Salient Weather Conditions & Forecast

Monsoon trough runs through Uttar Pradesh, Bihar, West Bengal to Assam across northern part of Bangladesh. Monsoon is fairly active over Bangladesh and weak to moderate over North Bay. As per the weather forecast received from the Bangladesh Meteorological Department (BMD) during next 24 hours, light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at a few places over the district. Day and night temperature may rise slightly over the district. As per the outlook issued by BMD for next 72 hours, rain/thunder showers is likely to increase at the end period. During last four days' light rainfall occurred and as per the quantitative medium range weather forecast, trace to light rainfall is likely during next five days. Detail advisories are given below

Aus Paddy:

Recovery to Panicle Initiation stages

- Maintain in 3-4 inches water level. Maintain 5-7 cm water level at panicle initiation stage.
- Inspect & monitor disease-pest e.g blast, rice bug, stem borers and others and take appropriate plant protection measures
- After decrease in water level from aus paddy field, apply potash fertilizer as top dress by filling gap after 5-7 days and also urea fertilizer after 3-5 days. Urea fertilizer should not be applied if the paddy is in early tillering stage. After 5-7 days, if necessary, urea fertilizer should be applied by breaking the tillers and filling the gap. At this time, it is better to use additional 5 kg of potash fertilizer per bigha in all types of aus paddy fields.
- If the population of Thrips and Green Leaf Hopper exceeds 25%, apply Malathion group insecticide at the rate of 1ml/1 litre of water.

Aman Paddy

- Advise for preparation of seeded for Aman paddy in *kharif season*.
- Start sowing of dry seeded long duration paddy after proper tillage operation like ploughing, weeding etc.
- Before sowing seed should be treated by Diethane M-45 properly
- Select high land for Aman rice seedbed where there is less probability of flooding. Raise seedlings by using floating or Dapog seedbed technology where raised land is not available.
- Apply 25-30 kg manure/decomposed cow dung in each decimal seedbed. In order to get good result, applying 2-3 kg vermicompost in same land size.
- Community based rice nursery may be raised to avoid loss of nursery due to heavy rainfall & flooding.

Vegetables:

- Hand pollination is effective for more production in case of tassel gourd and pointed gourd. Moreover, advised to cut aged leaves and continue others intercultural operations in vegetables land.
- Wash away sticky mud from base and stems of vegetable plants by spraying fresh water; to prevent rotting, spray Copper hydroxide or copper oxychloride @4g per litre of water. Staking of bamboo should be provided to crops like tomato, chilli and brinjal for better support.
- Arrange for adequate drainage to avoid water stagnation in the crop fields. Drain out excess water from the fields. Avoid irrigation, intercultural operation and application of plant protection measures and fertilizers to the standing crops. Harvest the matured crop immediately and store the harvested products at safe places.

Horticulture:

- During the month, fruit ripening will be started completely. Take special care like fertilization, and sticking and other intercultural operations for transplanted seedling/sapling.
- It is high time for transplanting fruit-tree (Mango, coconut, palm) tree, neem) plantation Arrange for fencing the transplanted seedlings/sapling for controlling cattle attract and keep the garden net and clean.
- During the month fruiting setting starts and banana bunch will grow.
- Besides, need to control banana beetle, panama and others diseases-pests including continue intercultural operations and has to inspect the field regularly.
- Weather is congenial for sigatoka leaf spot disease in banana; cut and remove all severely affected leaves and spray Propiconazol 1ml/ litre of water. Weather is congenial for burrowing nematode in banana; apply neem cake (1 kg/plant) at the time of planting to reduce the pest population. Store the harvested banana bunches in dry places to prevent disease incidence due to rainfall.
- Provide mechanical support to banana, young fruit plants and vegetables and carry out propping up in sugarcane to prevent the crops from lodging due to strong winds. Keep harvested produce at safer places. Keep cattle/goats indoor during thunderstorm/dust storm.
- Weather is congenial for bud rot disease in coconut; spray 1% Bordeaux mixture on the spindle leaf and crown of palms. Make provisions for small drains at required interval in the orchards to drain out excess water.

Jute:

- Foot rot, stem rot, and others diseases-pests may be infested and so has inspect the field regularly.
- Keep the jute land weed free by weeding and uproot comparatively under growth plant during the time.
- There is a chance of incidences of Semilooper. Spray Imidocholoprid/Chlorocyrin/ Nitro @ 1ml/litre of water. Semilooper and mites may infest in drought and humid weather and so placing 10-12 bamboo sticks/ plant branch @33 decimal land for controlling insects as environment friendly way. Besides, inspect the land regularly.
- It is now in vegetative stage. Monitor the pest & diseases incidences and take appropriate measures.

- To protect capsularis jute crop (>4 ft tall) from lodging due high wind speed, the peripheral jute plants (round the field) need to be tied together by taking 4-5 jute plants in a consecutive manner.
- There is a chance of incidences of Hairy Caterpillar. Following control measures is advised.
- Collect the egg mass and destroy
- Use light traps.
- Spray Imidocholoprid/Chlorocyryn/ Nitro @ 1ml/litre of water.
- There is a chance of incidences of Semiloper. Spray Imidocholoprid/Chlorocyryn/ Nitro @ 1ml/litre of water.

Livestock

- Keep animals indoor during heavy rainfall. For cow and buffalo HS and BS vaccine should be given. During Monsoon keep chicks & birds in a relatively warmer places or use electrical bulbs.
- Immunize the farm cattle against FMD (Foot & Mouth Disease), HS (Hemorrhagic Septicemia) and BQ (Black Quarter) diseases after consultation with local veterinary doctor.

Poultry

- For all age groups of poultry birds including local, broiler and layers give one dose of vaccine against Newcastle/Raniket disease in drinking water (Lasota/ F-1vaccine). Maintain free circulation of Air in poultry shed.

Fishery

- Repair damaged dykes of fisheries ponds and take care so that fishes do not escape. Maintain outlet pipe from the pond so that excess water pass away and the water level is maintain.
- Fish grower framers are advised to collect the healthy fingerlings from the authentic sources as there is enough water in the fish pond.
- Remove the unwanted fishes and predator by repeated netting before leaving the new fingerling in the fish pond.
- This is time for release of new batches of fishes in fish ponds.
- Select healthy, diseases free breeds of fishes from authentic sources after consulting with fishery officials. Before releasing, the fingerlings should be treated with Potassium permanganate (@ 3-5%.