



# NATIONAL AGROMETEOROLOGICAL ADVISORY SERVICE BULLETIN

10 November 2021 (WEDNESDAY)  
[FOR THE PERIOD 10.11.2021 – 14.11.2021]



## DISCLAIMER

Under the Agro-meteorological Information System Development Project, Agromet Advisory Service Bulletins at National & Sixty Four (64) districts have been prepared on experimental basis. These bulletins are being displayed through the website of Department of Agricultural Extension for valuable comments/ suggestions from all concerned.

Please Contact: Farhana Hoque, Sabuj Roy | E-mail: [adamisd@dae.gov.bd](mailto:adamisd@dae.gov.bd)  
Phone No.: 02-55'028414, 02-55028418

## ISSUED BY

Agro-Meteorological Information Systems  
Development Project  
Department of Agricultural Extension (DAE)  
Ministry of Agriculture  
In collaboration with Bangladesh  
Meteorological Department

## **Special Agromet Advisories for CORONA virus (COVID-19) transmission in Bangladesh**

### **Agromet Advisories**

Farmers and all other farm managements are advised to follow the guidelines of Government of Bangladesh to avoid infection and social transmission of CORONA virus (COVID-19). Precautions and safety measures should be taken up to prevent the Carona virus spread. Simple measures include social distancing, maintaining personal hygiene by washing hands with soap, wearing of face mask, drink hot water, stay at home and cleaning of implements and machinery. Farmers should not work in a group; consult with a doctor in case of any symptom. Please stay at home does not visit the field until it is very necessary.

### **Salient Weather Conditions & Forecast**

As per the weather forecast of Bangladesh Meteorological Department (BMD) during next 24 hours' Weather may remain dry over the country. Night temperature may rise slightly and day temperature may remain nearly unchanged over the country. During last four days, no rainfall occurs of the country and as per the quantitative medium range weather forecast is likely no rainfall in the next five days. Detail Agromet advisory are given below:

#### **Aman**

- Maintain the required water level in the land.
- In case of infestation of stem borer insects, apply 180-190 gm of Cartap group or 10 gm of Thiamethoxam + Chlorantraniliprol group of pesticides per bigha for control.
- In order to control Sheath Blight Disease, potash fertilizer should be divided into two equal installments, one part should be applied in the final cultivation of land preparation and the other part should be applied in combination with the last installment of urea fertilizer. Follow irrigation management in wet and dry manner periodically. Use fungicides in approved doses to successfully control the disease.
- In case of bacterial blight, in the early stages of the disease, mix 60 gm MOP, 60 gm thiovit in 10 liters of water and apply on 5 decimal of the land. If the disease occurs before the heading stage, 5 kg of potash fertilizer should be applied per bigha.
- In case of infestation of leaf-wrapping insects, apply 180-190 gm of Cartap group or 10 gm of Thiamethoxam + Chlorantraniliprole group of pesticides per bigha for control.
- Dew or drizzle rain is conducive to blast disease. In this type of weather, once in a while and 5-6 days after the emergence of heading, apply 54 gm of Trooper 75 WP / Difa 75 WP / Jill 75 WP or 33 gm of Nativo 75 WG or Tricyclazol / Strabin only per bigha of land. Mix well in liter of water and spray in the last afternoon.
- Rice Bug insect infestation may occur at the grain formation stage. Apply Carbaryl or Isoprocarb / MIPC group pesticides in approved doses if one Rice Bug insect is found in 2-3 bunches on an average.
- Use light traps to control insects.
- Drain out water from the paddy field 15 days before harvesting
- If the crop is 80% mature, collect it in a sunny day.

- Apply pesticides.

#### **Wheat:**

- Continue sowing wheat from 15th to 30th November.

#### **Potato:**

- November is the best time to plant potatoes. Continue planting potatoes in the current weather conditions. If there is no necessary moisture in the soil after sowing, provide irrigation as required.

#### **Maize:**

- Continue sowing the seeds. Apply first irrigation within 15-20 days of sowing, second irrigation within 30-35 days, third irrigation within 60-70 days and fourth irrigation within 85-89 days.

#### **Sugarcane**

- Sugarcane stems can be attacked by red rot disease. Infected trees should be removed from the land along with bushes as soon as the disease is seen. Water should be removed from the sugarcane field very quickly.
- To control the Sugarcane Stem Borer, remove the soil from the base of the sugarcane with a spade. Drain out accumulated water from the land. Old dried leaves should be removed from the trees, collected and burnt or buried under the ground. Egg parasitic hornet *Trichogramma Chilonis* should be released in the affected land at the rate of one gram per hectare per week (approximately 50,000). Apply Virtako 40 WG (Thiamethoxam + Chlorantraniliprole) to the infested sugarcane bush in the approved dose for control of stem borer insects with pesticides or Cartop granular pesticides such as Nokotap 6G cut the shallow drains on both sides of the sugarcane row and sprinkle in the drains then cover with soil or sprinkle at the base of the tree.
- Apply light irrigation and pesticides.

#### **Vegetables**

- Arrange for hand pollination of pumpkin vegetables. If beetles appear, kill them by hand in the morning and afternoon.
- In case of fly infestation in pumpkin, use pheromone and poison traps together. Pesticides of Alpha-Cypermethrin group can be used.
- In case of infestation of *eggplant* tip and fruit borer insects, cut and destroy the affected tip along with the insects. Insect reproduction is likely to be greatly reduced by using pheromone traps. Use spray of approval dose of insecticides, if necessary or recommended local organic insecticides may be used.
- In case of spider infestation, soak 1 kg half-broken neem seeds in 20 liters of water for 24 hours and spray the water (after chopping) on the underside of the leaves. If the

infestation is severe, spray the spider mite Omit 57 EC at the rate of 2.0 ml per liter of water or Vertimek 1.8 EC at the rate of 12 ml per 10 liters of water.

- Apply light irrigation and pesticides.

### **Horticultural crops**

- In case of spider infestation on the coconut tree, all the young caconut of the affected tree should be cut down and burnt. Infected trees and nearby young trees should be sprayed with any spider repellent such as Omite 57 EC mixed in 1.5 ml of one liter of water.
- In case of symptoms of Sigatoka disease on banana leaves, 0.5 ml of Score or 2 g of Noin or Bavistin or 0.1 ml of Econazole / Folico should be mixed in every liter of water and sprayed 2-3 times at intervals of 15-20 days.
- Apply Isoprocarb (MIPC) group pesticide if banana beetle infestation occurs.
- In case of infestation of milli bugs on papaya trees, the infected leaves / stems including insects should be collected and destroyed at the beginning of the attack. If the infestation is severe, mix 5 gms of soapy water or Admire 200 ml at the rate of 0.25 ml per liter of water and spray 2-3 times in 15 days.
- Apply light irrigation and pesticides.

### **Livestock:**

- Cattle should be fed with hay, raw grass and hand-made granular feed.
- Vaccinate cattle regularly to prevent disease.
- Sprinkle potassium permanganate solution to remove odors around.

### **Poultry:**

- Vaccinate chickens regularly to prevent disease.
- Sprinkle potassium permanganate solution to remove odors around.

### **Fishery**

- Prepare the pond for the next season by collecting the fish to be collected at the present time.
- Apply 200-250 grams of lime per decimal to clean the water of ponds where fish cannot be extracted.

## Realized Weather at different locations in the country

Rainfall for last 24 hours (till 06 AM on 10 November 2021) and maximum temperature on 09 November and minimum temperature on 10 November 2021 is given below:

Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)	Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)
Dhaka	Dhaka	00	31.6	19.5	Rajshahi	Rajshahi	00	30.8	18.0
	Tangail	00	31.2	17.0		Ishurdi	00	30.3	18.3
	Faridpur	00	31.0	18.2		Bogura	00	30.5	19.0
	Madaripur	00	30.2	18.4		Badalgachhi	00	30.3	16.6
	Gopalganj	00	30.4	17.4		Tarash	00	31.0	18.5
	Nikli	00	30.5	20.0					
Mymensingh	Mymensingh	00	31.3	18.0	Rangpur	Rangpur	00	31.0	18.6
	Netrokona	00	30.0	18.0		Dinajpur	00	31.5	17.2
						Sayedpur	00	32.3	17.5
Chattogram	Chattogram	00	32.5	23.5		Tetulia	00	32.0	<b>13.8</b>
	Sandwip	00	33.6	20.5		Dimla	00	31.0	17.0
	Sitakunda	xx	30.8	xx		Rajarhat	00	30.0	16.6
	Rangamati	00	32.5	22.0	Khulna	Khulna	00	31.5	19.4
	Cumilla	00	31.8	19.8		Mongla	00	31.4	20.0
	Chandpur	00	32.4	20.0		Satkhira	00	30.4	19.5
	M.Court	00	32.6	21.8		Jashore	00	32.0	17.4
	Feni	00	<b>34.0</b>	20.5		Chuadanga	00	31.2	17.2
	Hatiya	xx	32.4	xx	Kumarkhali	00	30.8	17.0	
	Cox's Bazar	00	33.3	24.6	Barishal	Barishal	00	31.8	18.5
Kutubdia	00	31.8	22.6	Patuakhali		00	32.0	19.7	
Teknaf	xx	30.8	xx	Khepupara		00	31.2	20.1	
Sylhet	Sylhet	00	31.9	18.3		Bhola	00	31.4	18.6
	Srimangal	00	32.3	19.2					

### Highlights:


- Country average bright sunshine hour was 8.32 hours per day during last week.
- Country average free water loss during the previous week was 3.11 mm per day.





### Weather Forecast Valid for 24 Hours commencing 06 PM Today:




**Forecast:** Weather may remain dry over the country.

**Temperature:** Night temperature may rise slightly and day temperature may remain nearly unchanged over the country.

**Spatial Distribution of weather parameters for the Week ending 09 November 2021**









### Weather Forecast and Advisory for the Period of 08.11.2021 to 14.11.2021


Bright sunshine hour is likely to be from 6.75 to 7.50 hours per day during this week. Average free water loss is likely to be from 3.25 to 4.25 mm per day during this week.

- Weather may remain mainly dry with temporary partly cloudy sky over the country during the 1st half and light rain (04-10 mm/d)/drizzle may occur at one or two places over Khulna, Barishal & chattogram divisions and weather may remain mainly dry elsewhere over the country during the 2nd half of the period.
- Light fog may occur over northern part & river basin of the country from early morning during the period.
- Day & night temperature may remain nearly unchanged during the 1st half and slight fall in day & night temperature is expected over the country during the 2nd half of the period.

### District Wise Quantitative Weather Forecast for next Five days from 10 Nov.-14 November 2021









## Different Satellite Products over Bangladesh


NOAA/VIIRS BLENDED NDVI composite for the week. No. 43 (22 October-28 October) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VCI composite for the week No. 43 (22 October-28 October) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED TCI composite for the week No. 43 (22 October-28 October) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VHI composite for the week No. 43 (22 October-28 October) over Agricultural regions of Bangladesh

