







Crop Weather Calendar of Maize: Dhaka Region (Districts: Dhaka, Tangail, Gazipur, Narsingdi, Narayanganj, Munshiganj, Manikgonj)
Bangladesh

Region:	Maize (Rabi)										Duration: 112 days					
Dhaka	October		November				December				January				February	
Months	October		November				December				January				February	
Std.Week/Normal	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6
Rainfall (mm)	26.0	12.5	10.0	6.0	1.5	4.5	1.5	4.5	1.0	3.5	1.0	0.0	0.0	0.5	3.5	3.5
Max. Temp. (°C)	31.5	30.9	30.6	29.8	29.0	28.3	27.6	26.2	25.5	24.8	24.0	23.9	24.5	25.0	26.2	27.2
Min. Temp. (°C)	22.3	21.5	20.5	18.7	17.7	16.1	15.5	14.7	13.5	12.9	12.4	12.0	12.0	12.1	13.6	14.3
Mean Temp. (°C)	26.9	26.2	25.5	24.3	23.3	22.2	21.5	20.5	19.5	18.8	18.2	17.9	18.3	18.6	19.9	20.7
RHmax (%)	94.8	94.5	95.0	94.7	94.6	94.3	94.8	95.3	95.1	95.0	94.6	94.3	93.8	93.4	92.4	92.1
RHMin (%)	55.5	54.6	51.5	49.3	47.5	46.3	47.6	49.4	49.8	49.1	50.3	48.8	46.2	43.3	42.3	39.4
RHmean (%)	75.2	74.6	73.2	72.0	71.0	70.3	71.2	72.3	72.5	72.1	72.4	71.6	70.0	68.3	67.3	65.7
SShr (hrs)	54.0	49.5	51.5	51.5	52.0	51.5	48.5	46.5	44.5	44.5	42.5	42.5	45.0	49.0	49.5	51.5
WD (deg)	190	190	189	199	212	211	219	224	225	233	238	238	232	242	241	236
WS(Km/hr)	3.8	3.7	3.7	3.7	3.7	3.6	3.7	3.7	3.7	3.7	4.1	4.0	4.1	4.3	4.3	4.5

	 					
	sowing	Germination	Vegetative Growth	Silking & Tasseling	Cob Formation	Maturity to Harvesting

Favorable Weather Conditions					
Temperature	At least 10° C for germination			22-25° C	23-27° C
Light Intensity		≤200% of normal			
Relative Humidity		High			
Soil Temperature		Above 16° C			
Normal phase wise water requirement (mm)	76	120	190	145	100

Congenial Weather Condition for Pests & Diseases	
Fall Army worm	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation and adult emergence. Emergence in sandy-clay and clay-sand soils was directly proportional to temperature and

	inversely proportional to humidity. Above 30°C the wings of adults tend to be deformed. Pupae require a threshold temperature of 14.6°C and 138 day-degrees C to complete their development.			
Seed Rot	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation.			

Weather Warning					
Rain		>50 mm/day	>100 mm/day	>50 mm/day	
Duration of wet spell		>25 mm for 3 days	>50 mm for 4 days	20 mm for 4 days	
Cloudy weather			Cloudy weather	Cloudy weather	
High wind		>50 km/hr	>40 km/hr	>30 km/hr	
Temperature		Minimum Temperature <10 ° C	Minimum Temperature <10° C	Minimum Temperature <10 ° C	