







Crop Weather Calendar of Maize: Dhaka Region (Districts: Dhaka, Tangail, Gazipur, Narsingdi, Narayanganj, Munshiganj, Manikgonj)
Bangladesh

Region:	Maize (kharif-1)														Duration: 112 days	
Dhaka																
Months	April		May				June				July				August	
Std.Week/Normal	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Rainfall (mm)	31.5	42.5	46.0	55.5	55.0	75.5	70.0	72.5	86.5	81.0	77.5	82.5	73.5	91.0	81.5	67.0
Max. Temp. (°C)	34.1	33.8	33.6	33.7	33.7	33.2	33.5	33.4	32.8	32.2	32.2	32.1	32.0	31.7	32.2	32.2
Min. Temp. (°C)	23.7	23.6	23.7	24.2	24.6	24.9	25.4	25.8	25.9	26.2	26.2	26.1	26.2	26.0	26.0	26.2
Mean Temp. (°C)	28.9	28.7	28.6	28.9	29.2	29.0	29.4	29.6	29.3	29.2	29.2	29.1	29.1	28.9	29.1	29.2
RHmax (%)	91.4	91.7	92.3	92.8	92.9	93.5	93.8	94.0	94.5	94.4	94.7	94.7	94.1	94.5	94.3	94.0
RHMin (%)	52.0	55.1	56.7	58.6	60.4	63.5	65.0	66.7	69.9	71.5	72.9	73.5	73.1	73.5	71.1	71.0
RHmean (%)	71.7	73.4	74.5	75.7	76.7	78.5	79.4	80.3	82.2	83.0	83.8	84.1	83.6	84.0	82.7	82.5
SShr (hrs)	55.0	54.5	51.5	48.0	48.0	42.5	42.0	38.5	31.0	26.0	26.0	27.0	27.5	28.5	32.0	33.5
WD (deg)	167	156	159	164	156	155	157	161	157	157	158	162	157	161	159	156
WS(Km/hr)	6.6	6.7	6.8	6.5	6.4	6.1	5.8	5.6	6.3	6.5	6.5	5.8	5.9	6.1	5.9	6.1

						
	sowing	Germination	Vegetative Growth	Silking & Tasseling	Cob Formation	Maturity to Harvesting

Favorable Weather Conditions					
Temperature	At least 10° C for germination			22-25° C	23-27° C
Light Intensity		≤200% of normal			
Relative Humidity		High			
Soil Temperature		Above 16° C			
Normal phase wise water requirement (mm)	76	120	190	145	100

Congenial Weather Condition for Pests & Diseases

Fall Army worm	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation and adult emergence. Emergence in sandy-clay and clay-sand soils was directly proportional to temperature and inversely proportional to humidity. Above 30°C the wings of adults tend to be deformed. Pupae require a threshold temperature of 14.6°C and 138 day-degrees C to complete their development.			
Seed Rot	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation.			

Weather Warning					
Rain		>50 mm/day	>100 mm/day	>50 mm/day	
Duration of wet spell		>25 mm for 3 days	>50 mm for 4 days	20 mm for 4 days	
Cloudy weather			Cloudy weather	Cloudy weather	
High wind		>50 km/hr	>40 km/hr	>30 km/hr	
Temperature		Minimum Temperature <10 °C	Minimum Temperature <10° C	Minimum Temperature <10 °C	