







Crop Weather Calendar of Maize: Mymensingh Region (Districts: Mymensingh, Kishoregonj, Jamalpur, Netrokona, Sherpur) Bangladesh

Region: Mymensingh	Maize (Rabi)												Duration: 112 days			
Months	December				January				February				March			
Std.Week/Normal	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12
Rainfall (mm)	1.0	3.0	0.0	2.0	0.0	3.0	1.0	0.0	2.0	1.0	5.0	7.0	1.0	4.0	6.0	11.0
Max. Temp. (°C)	27.6	26.3	25.4	25.0	24.4	24.0	24.0	25.1	25.9	26.6	27.4	28.1	29.1	30.3	31.0	31.9
Min. Temp. (°C)	15.0	14.0	13.3	12.4	12.0	11.6	11.9	12.0	13.0	13.7	15.0	15.9	16.6	17.4	18.6	19.9
Mean Temp. (°C)	21.3	20.2	19.4	18.7	18.2	17.8	18.0	18.6	19.5	20.2	21.2	22.0	22.9	23.9	24.8	25.9
RHmax (%)	95.3	95.6	95.6	95.3	95.1	94.4	94.4	93.4	92.6	92.1	91.4	90.6	90.9	90.6	90.1	89.7
RHMin (%)	52.3	54.1	53.6	53.7	53.6	53.1	51.4	47	47.9	45.3	45	45.1	44.3	43.6	44.6	48.3
RHmean (%)	73.8	74.9	74.6	74.5	74.4	73.8	72.9	70.2	70.3	68.7	68.2	67.9	67.6	67.1	67.4	69.0
SShr (hrs)	49.0	43.0	40.0	42.0	41.0	39.0	41.0	45.0	44.0	48.0	50.0	50.0	51.0	53.0	53.0	48.0
WD (deg)	191	206	199	221	230	236	216	238	228	218	222	210	214	202	208	179
WS(Km/hr)	3.0	3.5	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.9	4.4	5.4	5.0	5.4	5.6	6.3
																
	sowing	Germination	Vegetative Growth						Silking & Tasseling			Cob Formation			Maturity to Harvesting	
Favorable Weather Conditions																
Temperature	At least 10°C for germination												22-25°C		23-27°C	
Light Intensity			≤200% of normal													
Relative Humidity			High													
Soil Temperature			Above 16°C													
Normal phase wise water requirement (mm)	76		120						190				145		100	
Congenial Weather Condition for Pests & Diseases																
Fall Army worm	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation and adult emergence. Emergence in sandy-clay and clay-sand soils was directly proportional to temperature and inversely proportional to humidity. Above 30°C the wings of adults tend to be deformed. Pupae require a threshold temperature of 14.6°C and 138 day-degrees C to complete their development.															
Seed Rot	A threshold temperature of 10.9°C and 559 day-degrees C is required for development. Sandy-clay or clay-sand soils are suitable for pupation.															
Weather Warning																
Rain	>50 mm/day						>100 mm/day				>50 mm/day					
Duration of wet spell	>25 mm for 3 days						>50 mm for 4 days				20 mm for 4 days					
Cloudy weather							Cloudy weather				Cloudy weather					
High wind	>50 km/hr						>40 km/hr				>30 km/hr					
Temperature	Minimum Temperature <10°C						Minimum Temperature <10°C				Minimum Temperature <10°C					