Crop Weather Calendar of RobiTomato:Dinajpur Region (Districts:Dinajpur, Thakurgaon, Panchagaor) Bangladesh

Dinajpur Region				i Toma			Life span 120-130											
Months	November						Decer	nber		1	Janu		February					
Std.Week	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Rainfall (mm)	5.0	3.0	0.0	0.0	0.0	0.0	3.0	0.0	4.0	2.0	2.0	3.0	2.0	1.0	2.0	3.0	1.0	
Max. Temp. (°C)	30.3	29.7	29.0	28.7	28.0	27.0	25.4	24.9	23.9	22.6	22.4	23.0	23.7	25.0	25.9	26.7	27.7	
Min. Temp. (°C)	19.4	18.3	16.9	15.6	14.0	13.3	12.9	12.0	10.9	10.7	10.3	10.0	10.6	11.1	12.1	13.6	14.1	
Mean Temp. (°C)	24.9	24.0	23.0	22.2	21.0	20.2	19.2	18.5	17.4	16.7	16.4	16.5	17.2	18.1	19.0	20.2	20.9	
RHmax (%)	96.0	95.9	95.9	95.3	95.6	96.1	96.4	96.3	96.6	96.7	96.3	96.4	95.9	94.7	94.0	93.4	92.0	
RHMin (%)	55.4	53.0	51.6	48.1	47.6	49.0	52.9	52.7	54.0	55.1	53.7	52.4	50.4	47.7	43.1	42.4	40.9	
RHmean (%)	75.7	74.5	73.7	71.7	71.6	72.5	74.6	74.5	75.3	75.9	75.0	74.4	73.1	71.2	68.5	67.9	66.4	
SShr (hrs)	55	56	52.0	54.0	53.0	48.0	42.0	41.0	37.0	34.0	34.0	37.0	41.0	44.0	46.0	52.0	54.0	
WD (deg)	149	144	154	164	177	183	193	200	213	225	238	243	227	241	225	230	225	
WS(Km/hr)	3.5	3.7	3.7	3.7	3.5	2.6	3.7	2.9	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
						4 9							P	5		X		
Stages	Germination and Vege Seedling Planting					Vegetative	ive Growth Flowering & Fruiting					Harvesting						
Favourable Weather																		
Temperature		18-25 °C																
Relative humidity										50-70%								
Soil temperature										20-25 °C								
Sunshine hour									1	6-8 hr							ı	
Favourable weather for dis																		
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																	
Tomato fruit borer		1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %																
Leaf miner		Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population															itive and significant	
Early blight					Dis	ease develo	ps at mode	erate to wa	arm (15°C					90% or gre	eater, mod	lerately to	heavy rainfall	
Late blight						Temper	rature: 15-	27°C, pro	longed ho	t dry days	increased d	isease infes	station; re	lative hum	nidity 75 to	o 90% or g	greater	
Fungal Wilting			Dry at	mospheri	c conditions	s, Max. Soil	temperatu	ire 35°C to	o 43.5°C;	Mini. Soil	temperature	e 25°C to 3	80°C; RH	: 30%-509	%			
Weather warning																		
Rainfall (mm)					≥20 r	nm rainfall	per day at	the end of			eptember-N	ovember)						
Cloudy weather		Cloudy condition																
Drought		Drought (moisture stress) due to excessive dry weather																
Duration of staying wet		Land remain waterlogged for 2 to 4 days																
Hailstorm					Late sea	son rainfall	with hails	torm										
Windspeed										15 km/hr								
Temperature (°C)									3	0 to 32°C								

Crop Weather Calendar of Kharif Tomato: Dinajpur Region (Districts:Dinajpur, Thakurgaon, Panchagaor) Bangladesh

Dinajpur Region				Tomato	(= 10 011		Life span 120-130											
Months		March						April				lay		June				
Std.Week	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
Rainfall (mm)	1.0	1.0	4.0	2.0	7.0	8.0	8.0	25.0	23.0	31.0	36.0	57.0	55.0	73.0	57.0	77.0	111.0	
Max. Temp. (°C)	28.6	30.1	31.1	32.0	32.1	33.0	33.4	33.0	32.4	32.3	32.9	33.0	33.0	33.4	33.3	33.0	32.3	
Min. Temp. (°C)	15.0	16.1	17.1	18.6	19.7	20.4	21.1	22.0	22.0	22.3	22.9	23.6	24.0	24.4	25.0	25.1	25.6	
Mean Temp. (°C)	21.8	23.1	24.1	25.3	25.9	26.7	17.3	27.5	27.2	27.3	27.9	28.3	28.5	28.9	29.2	29.1	29.0	
RHmax (%)	90.6	88.9	87.6	86.0	87.1	86.6	87.4	89.4	91.1	92.3	92.0	92.1	92.4	92.7	93.4	93.9	94.7	
RHMin (%)	38.4	35.3	35.1	36.4	43.3	40.9	43.4	50.9	53.7	57.6	57.3	59.3	62.4	62.3	65.4	67.7	71.4	
RHmean (%)	64.5	62.1	61.4	61.2	65.2	63.8	65.4	70.2	72.4	75.0	74.7	75.7	77.4	77.5	79.4	80.8	83.1	
SShr (hrs)	57.0	59.0	59.0	55.0	49.0	52.0	55.0	49.0	50.0	49.0	51.0	48.0	44.0	42.0	40.0	35.0	27.0	
WD (deg)	228	230	212	192	170	171	150	142	123	118	115	119	108	117	111	118	120	
WS(Km/hr)	3.8	4.2	5.4	5.6	5.0	5.6	4.8	4.8	5.4	3.9	3.7	4.3	3.7	3.9	3.7	3.7	3.7	
2						a de la companya de l							P	Y			*	
Stages		Germination and Veg Seedling Planting						tative Growth Flow			vering & Fruiting			Harvesting				
Favourable Weather																		
Temperature									26-32°C									
Relative humidity		50-80%																
Soil temperature		22-26 °C																
Sunshine hour									8-10 hr									
Favourable weather for o																		
White fly	15°C to	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																
Tomato fruit borer		1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61																
Leaf miner		Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population															itive and	
Early blight					Di	sease devel	ops at mode	rate to war						or greater	r, moderate	ly to heav	y rainfall	
Late blight							erature: 15-2						tion; relativ	e humidit	y 75 to 90%	or greate	r	
Fungal Wilting	Dry at	mospheric	conditions	, Max. Soil	temperati	ire 35°C to	43.5°C; Mii	ni. Soil ten	nperature 2	25°C to 30	°C; RH: 30	9%-50%						
Weather warning																		
Rainfall (mm)			≥20 n	nm rainfall	per day at	the end of	he rainy sea											
Cloudy weather					ı			Clo	oudy condi	tion								
Drought							10 -				Droug	ht (moisture	e stress) du	e to excess	ive dry we	ather		
Duration of staying wet							ogged for 2											
Hailstorm					Late	season rain	fall with hai		. 151 "									
Windspeed									≥15 km/hi									
Temperature (°C)									30 to 32°C	,								