Crop Weather Calendar of Rabi Chili(Anthracnose disease): Rajshahi Region (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

	Rajshahi Region						Life span 170-180 days																
	Months		N	ovembe	r			Dece	ember			Jan	uary			Feb	ruary				March		
Weather warning	Max. Temp. (°C)			28 °C to 32 °C for disease development																			
	Min. Temp. (°C)									22 °C 1	to 25 °C	for dis	ease de	velopn	nent								
	RH Max (%)							Initial	infecti	on requ	ired 95	% or ab	ove rela	ative h	umidity	for 24 l	hrs.						
> 5	RH Min (%)											809	6										
	Rainfall (mm)		Frequent rainy and warm weather alone with prolonged leaf wetness favored disease development																				
	Std.Week/Normal	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
normal ther	Rainfall (mm)	7.0	5.0	1.0	0.0	2.0	1.0	3.0	0.0	4.0	1.0	1.0	2.0	1.0	4.0	3.0	2.0	3.0	5.0	4.0	3.0	4.0	14.0
ori	Max. Temp. (°C)	31.7	31.0	30.4	30.4	29.6	27.0							24.9			28.1	29.3		32.1	33.1	34.6	35.0
	Min. Temp. (°C)	23.1	21.4			17.6	14.0		12.4									14.1		16.4	17.6	19.4	20.6
eekly wea	RHmax (%)	96.1	96.0	95.9	96.1	95.7	94.7	95.9	95.3	95.7	95.6	95.1	95.3	95.3	95.3	94.0	93.9		91.6	90.3	88.7	88.3	89.1
We	RHMin (%)	60.9	57.4	55.9	51.9	49.4	46.9	50.1	50.0	50.0	50.1	48.4	47.9	45.7	44.7	42.0	40.3	36.6	34.6	32.7	30.4	30.4	35.9
	SShr (hrs)	53.0	56.0	55.0	57.0	54.0	51.0	47.0	46.0	44.0	42.0	42.0	46.0	52.0	51.0	55.0	56.0	58.0	59.0	59.0	59.0	59.0	52.0
	Stage	عے	T.					溪															
		22 °C to 25 °C for disease development Initial infection required 95% or above relative humidity for 24 hrs. 80%	esting																				
	Disease cycle			0	O NAME OF THE PARTY OF THE PART	e										Feb. 3.0 dity for 24 h 6.0 3.0 5.6 26.7 2.0 12.3 5.3 94.0 55.0 55.0							
											After occurrence of disease the disease cycle continued up to fruiting stage an								and ha	rvesting	and harvesting stage		

Crop Weather Calendar of Kharif Chili(Anthracnose disease): Rajshahi Region (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

	Rajshahi Region	Kharif season Chilli: Anthracnose disease									Life span 170-180 days												
Weather warning	Months	April may June July August																					
	Max. Temp. (°C)	28 °C to 32 °C for disease development																					
	Min. Temp. (°C)	22 °C to 25 °C for disease development																					
	RH Max (%)	Initial infection required 95% or above relative humidity for 24 hrs.																					
	RH Min (%)	80%																					
	Rainfall (mm)	Frequent rainy and warm weather alone with prolonged leaf wetness favored disease development																					
Weekly normal weather	Std.Week/Normal	78	%	<i>></i> ७	১৭	74	79	২০	২১	২২	২৩	২৪	২৫	২৬	২৭	২৮	২৯	೨೦	৩১	৩২	೨೨	৩৪	৩৫
	Rainfall (mm)	٥.0د	٥.٥	٥.هد	٥.٩٧	২৪.০	২৯.০	లల .0	৩৬.০	৩ 8.0	8b.o	65. 0	१४.०	৭২.০	৬৬.০	93.0	96.0	৭২.০	৫২.০	৫৭.০	৬8.০	৫৩.০	৫٩. ٥
	Max. Temp. (°C)	৩৬.১	৩৬.৭	৩৬.৪	৩৫.৯	৩৫.৪	৩৫.৩	৩৫.১	৩৫.০	৩৫.৩	৩৫.০	৩৩.৯	৩৩.১	೨೨.೦	৩২.৭	৩২.৯	৩২.৩	৩২.৬	৩২.৯	৩২.৯	৩২.৭	೨೨.೦	೨೨.೦
	Min. Temp. (°C)	২২.১	২৩.০	২৩.৪	২৩.৯	২৩.৭	২৪.৩	২৪.৬	২৫.০	২৫.৪	২৫.৯	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬১	২৬.০	২৬.১	২৬.০
	RHmax (%)	৮৮.৯	৮৯.০	৯০.১	৯০.৭	৯২.১	৯২.৭	৯৩.৩	৯৩.১	৯৩.৭	৯৪.৬	৯৫.৪	৯৬.১	৯৬.৬	৯৬.৬	৯৬.৯	৯৭.০	৯৬.৯	৯৬.৭	৯৬.৬	৯৬.৭	৯৬.১	৯৬.৪
	RHMin (%)	೦8.೦	৩৪.৬	8.48	86.5	¢0.0	৫১.৯	৫৫.৬	৫৮.৩	৫৮.৯	৬২.৬	৬৭.৯	۹۵.۵	৭৩.৪	৭৩.৭	৭৪.৩	৭৫.৩	৭৩.৯	૧૨.৬	૧૨.৬	48.\$	૧૨.૦	৭০.৯
	SShr (hrs)	¢9.0	৫৭.০	৫٩. ٥	৫৬.০	৫২.০	৫৩.০	0.83	8b.o	65.0	8b.o	৩৫.০	ు .ం	ు .ం	೨೦.೦	೨೦.೦	೨೦.೦	೨8.0	೨8.0	৩৬.০	ు ం.ం	೦8.೦	80.0
Stage Disease cycle																							
		Spouting & Seedling vegetative of						growth Flowerin					ng & fruiting						Harvesting				
			d e																				
			Spore Spore penetration							Disease development After occurrence of disease the disease cycle continued up to fruiting stage and harvesting stage										gstage			