



































Crop Insect Weather Calendar of *Rabi* season Brinjal Shoot and Fruit Borer (BSFB): Rajshahi Region (Districts: Rajshahi, Chapainawabgonj, Naogaon, Natore) Bangladesh

Region: Rajshahi		<i>Rabi</i> Season Brinjal: Brinjal Shoot and Fruit Borer (BSFB)											Life span 140-150 day											
Weather warning	Months	November			December				January				February				March							
	Max. Temp. (°C)	33.2																						
	Min. Temp. (°C)	10.7																						
	RH Max (%)	96.6																						
	RH Min (%)	35.4																						
	Rainfall (mm)	39.5																						
Weekly normal weather	Std. Week/Normal	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12		
	Rainfall (mm)	7.0	5.0	1.0	0.0	2.0	1.0	3.0	0.0	4.0	1.0	1.0	2.0	1.0	4.0	3.0	2.0	3.0	5.0	4.0	3.0	4.0		
	Max. Temp. (°C)	31.7	31.0	30.4	30.4	29.6	27.0	26.0	25.4	24.4	24.0	23.9	24.0	24.9	25.6	26.7	28.1	29.3	30.4	32.1	33.1	34.6		
	Min. Temp. (°C)	23.1	21.4	20.7	19.3	17.6	14.0	13.1	12.4	11.6	11.0	10.9	10.9	10.9	12.0	12.3	13.6	14.1	15.1	16.4	17.6	19.4		
	RH max (%)	96.1	96.0	95.9	96.1	95.7	94.7	95.9	95.3	95.7	95.6	95.1	95.3	95.3	95.3	94.0	93.9	92.0	91.6	90.3	88.7	88.3		
	RH Min (%)	60.9	57.4	55.9	51.9	49.4	46.9	50.1	50.0	50.0	50.1	48.4	47.9	45.7	44.7	42.0	40.3	36.6	34.6	32.7	30.4	30.4		
	SS hr (hrs)	53.0	56.0	55.0	57.0	54.0	51.0	47.0	46.0	44.0	42.0	42.0	46.0	52.0	51.0	55.0	56.0	58.0	59.0	59.0	59.0	59.0		
Stages																								
	Germination and Seedling Planting		Vegetative Growth				Flowering			Fruiting			Harvesting											
Insect Life cycle																								
	Eggs		Larva		Pupa		Adult		Eggs		Larva		Pupa		Adult		Eggs		Larva		Pupa		Adult	

Crop Insect Weather Calendar of *Kharif* season Brinjal Shoot and Fruit Borer (BSFB): Rajshahi Region (Districts: Rajshahi, Chapainawabgonj, Naogaon, Natore) Bangladesh

Region: Rajshahi		Kharif Season Brinjal: Brinjal Shoot and Fruit Borer (BSFB)											Life span 140-150 day													
Weather warning	Months	May					June					July					August					September				
	Max. Temp. (°C)	33.2																								
	Min. Temp. (°C)	10.7																								
	RH Max (%)	96.6																								
	RH Min (%)	35.4																								
	Rainfall (mm)	39.5																								
Weekly normal weather	Std. Week/Normal	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39			
	Rainfall (mm)	102.5	107.5	122.0	116.5	148.5	152.0	168.0	129.5	137.5	130.0	110.0	129.5	129.0	123.5	103.0	118.5	105.0	96.0	107.0	94.0	82.0	81.5			
	Max. Temp. (°C)	31.5	31.6	31.9	32.2	31.5	31.9	31.6	31.9	31.9	31.7	32.0	31.8	32.2	32.3	32.8	32.2	32.2	32.5	32.2	32.2	32.2	31.7			
	Min. Temp. (°C)	22.0	22.7	23.2	23.6	24.0	24.4	24.9	25.0	25.0	25.0	25.2	25.0	25.0	25.1	25.3	25.2	25.0	25.0	25.0	25.0	25.0	24.2			
	RH max (%)	94.0	94.6	94.7	94.9	95.2	95.8	96.0	96.2	96.0	95.9	96.0	96.1	96.2	96.4	96.2	96.4	96.3	96.3	96.4	96.4	96.1	96.5			
	RH Min (%)	63.1	64.2	65.6	67.2	70.6	71.8	73.6	73.7	74.0	73.9	73.4	74.2	72.6	72.4	70.9	72.8	73.0	72.4	72.3	71.7	71.4	71.8			
	SS hr (hrs)	45.0	43.5	43.5	38.5	33.5	31.5	26.0	27.5	23.5	25.0	27.0	26.0	30	33.0	33.5	29.5	32.5	33.5	32.5	33.0	34.0	34.5			
Stages																										
	Germination and Seedling Planting		Vegetative Growth					Flowering					Fruiting			Harvesting										
Insect Life cycle																										
			Eggs	Larva	Pupa	Adult	Eggs	Larva	Pupa	Adult	Eggs	Larva	Pupa	Adult												