

		
		
<p><b>Agro-Meteorological Information Systems Development Project</b>  <b>Component-C of BWCSR</b>  <b>Department of Agricultural Extension</b></p>		

## [Special Agromet Advisories for Rainfall \(Updated\)](#)

**Issue Date: 14 March 2023**

**Updated issue date: 19 March 2023**

According to the information received from the Bangladesh Meteorological Department, light to heavy rainfall is likely to occur in all districts of Bangladesh from March 17 to 22. At this time, there may be thunderstorms somewhere. Based on the analysis of the data received from the Bangladesh Meteorological Department, the following advisories are given for all districts:

- Avoid irrigation, fertilizer and pesticide application.
- Harvest mature crops quickly and store in a dry and safe place.
- Boro rice may be infested by blast disease. Tricyclazole group fungicides like Trooper (50 gm of fungicide mixed well in 64 liters of water per 1 bigha land) or ezoxytrobino group fungicides like Nativo (40 gm fungicide mixed well in 64 liters of water per 1 bigha land) should be sprayed twice at an interval of 7-10 days.
- Bacterial Leaf Blight may occur in Boro rice. To protect the crop from this disease, 5 kg of potash fertilizer and 3.5 kg of gypsum per bigha should be applied after the rains stop. However, if the paddy plant has passed the panicle initiation stage, 60 grams of potash fertilizer, 60 grams of thiovit and 20 grams of zinc should be mixed well in 10 liters of water and sprayed on 5 percent of the land. Stop application of urea fertilizer after stormy rains.
- Keep drains clear so that water does not accumulate on the ground.
- Arrange for elevated bunding around and inside the boro paddy.
- Provide mechanical support to the standing banana, sugarcane and other horticultural crops.
- If the field condition is suitable after rainfall, then start sowing of jute seeds.



(Dr. Md. Shah Kamal Khan)  
 Project Director  
 Cell no. +8801712184274  
 e-mail: [kamalmoa@gmail.com](mailto:kamalmoa@gmail.com)